PRODUCT INFORMATION



PAF Acetylhydrolase (human) Polyclonal Antibody

Item No. 160603

Overview and Properties

Contonto	This vial contains 500 ul of pontido offinity purified polyclopal antibody
Synonyme:	$I_{\rm m}$ DIA DAE AH Distributing Easter Decenter Acetulbudy
Synonyms.	LP-PLA ₂ , PAP-AH, Platelet-activating factor Receptor Acetylhydrolase
Immunogen:	Synthetic peptide corresponding to the C-terminal region of human PAF-AH
Cross Reactivity:	(+) PAF-AH
Species Reactivity:	(+) Human; (-) Mouse
Uniprot No.:	Q13093
Form:	Liquid
Storage:	-20°C (as supplied)
Stability:	≥3 years
Storage Buffer:	TBS, pH 7.4, with 50% glycerol, 0.1% BSA, and 0.02% sodium azide
Host:	Rabbit
Application:	Western blot (WB); the recommended starting dilution is 1:200. Other applications were not tested, therefore optimal working concentration/dilution should be determined empirically.

Image



Lane 1: LDL fraction from human plasma Lane 2: Recombinant human PAF-AH (20 ng)

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/16/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Platelet-activating factor acetylhydrolase (PAF-AH) is a calcium-independent member of the phospholipase A₂ (PLA₂) family that hydrolyzes the acetyl group at the *sn*-2 position of PAF, a phospholipid mediator involved in a variety of biological activities.^{1,2} There are three types of PAF-AH: two intracellular forms, PAF-AH1 and PAF-AH2, which are ubiquitously expressed, and an extracellular form, lipoprotein-associated phospholipase A₂ (Lp-PLA₂), which is found in the plasma in association with LDL and HDL.³⁻⁵. Unlike PAF-AH1, which is largely restricted to hydrolysis of PAF, PAF-AH2 and Lp-PLA₂ exhibit broader substrate specificity, hydrolyzing phospholipids with oxidized or short *sn*-2 acyl chains in addition to PAF.^{2,3} PAF-AH1 has roles in neuronal development, protein trafficking, and spermatogenesis, and PAF-AH2 and Lp-PLA₂ have a protective roles against oxidative stress and atherosclerosis, respectively.⁶ Cayman's PAF-acetylhydrolase (human) Polyclonal Antibody can be used for Western blot (WB) applications.

References

- 1. Kono, N. and Arai, H. Platelet-activating factor acetylhydrolases: An overview and update. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* **1864(6)**, 922-931 (2019).
- McIntyre, T.M., Prescott, S.M., and Stafforini, D.M. The emerging roles of PAF acetylhydrolase. J. Lipid Res. 50(Suppl), S255-S259 (2009).
- Kono, N. and Arai, H. Intracellular platelet-activating factor acetylhydrolase, type II: A unique cellular phospholipase A₂ that hydrolyzes oxidatively modified phospholipids. *Enzymes* 38, 43-54 (2015).
- 4. Murakami, M. Lipoquality control by phospholipase A2 enzymes. Proc. Jpn. Acad. Ser. B Phys. Biol. Sci. 93(9), 677-702 (2017).
- Karasawa, K., Harada, A., Satoh, N., et al. Plasma platelet activating factor-acetylhydrolase (PAF-AH). Prog. Lipid Res. 42(2), 93-114 (2003).
- 6. Ashraf, M.A. and Nookala, V. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing (2021). Available from: https://www.ncbi.nlm.nih.gov/books/NBK557392/

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM