# **PRODUCT** INFORMATION



## 2-Aminopurine (hydrochloride)

Item No. 18609

CAS Registry No.: Formal Name:	1428126-74-2 9H-purin-2-amine, dihydrochloride	
Svnonvm:	2-AP	
MF:	$C_5H_5N_5 \bullet 2HCI$	H <sub>2</sub> N N N
FW:	208.1	
Purity:	≥98%	N.
Ex./Em. Max:	320/381 nm	H N
Supplied as:	A crystalline solid	• 2HCI
Storage:	-20°C	
Stability:	≥4 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

#### Laboratory Procedures

2-Aminopurine is supplied as a crystalline solid. Aqueous solutions of 2-Aminopurine can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of 2-Aminopurine in PBS, pH 7.2, is approximately 5 mg/ml. We do not recommend storing the aqueous solution for more than one day.

#### Description

2-Aminopurine is a fluorescent analog of guanosine and adenosine that is used as a site-specific probe of nucleic acid structure and dynamics.<sup>1</sup> It base pairs with cytosine in a wobble configuration or with thymine in a Watson-Crick geometry.<sup>1</sup> In addition to its usefulness in studies of base stacking interactions, 2-aminopurine has been used as a probe for DNA base flipping by methyl transferases.<sup>2</sup> 2-Aminopurine exhibits an excitation/emission of 320/381 nm, respectively, when adjusted to reduce interference with DNA base absorption as well as tryptophan fluorescence.<sup>2</sup> 2-Aminopurine inhibits the double-stranded RNA-dependent protein kinase, protein kinase R, whose activity mediates antiviral defense and participates in toll-like receptor signaling.<sup>3</sup>

#### References

- 1. Jean, J.M., and Hall, K.B. 2-Aminopurine fluorescence quenching and lifetimes: Role of base stacking. Proc. Nat. Acad. Sci. USA 98(1), 37-41 (2001).
- 2. Holz, B., Klimasauskas, S., Serva, S., et al. 2-Aminopurine as a fluorescent probe for DNA base flipping by methyltransferases. Nucleic Acids Res. 26(4), 1076-1083 (1998).
- 3. Endoh, Y., Chung, Y.M., Clark, I.A., et al. IL-10-dependent S100A8 gene induction in monocytes/ macrophages by double-stranded RNA. J. Immunol. 182(4), 2258-2268 (2009).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/26/2022

### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM